

What should I know before taking CS 30

- **Boolean Variables.** Boolean variables or simply Booleans are the most basic unit of data: the *bit*. The bit takes two values: 1 or “True”, and 0 or “False”.
- **Numbers and Arithmetic.** There are many types of numbers: *natural/whole numbers* $\{0, 1, 2, 3, \dots\}$, *integers* $\{\dots, -2, -1, 0, 1, 2, \dots\}$, which also include negative numbers, *rational numbers*, which are of the form p/q where both p and q are integers, and *real numbers* which can be represented on the number line can be approximately represented by *decimals*.
- **Arithmetic.** Any two numbers can be added, subtracted, multiplied, and divided. Rationals are *closed* under these *operations*, that is, the sum/difference/product/ratio of any two rationals is rational. This is *not true* for integers. For instance, 2 divided by 4 is not an integer.
- **Associativity and Commutativity.** Addition and Multiplication are *associative* and *commutative*. That is, $(a + b) + c$ is the same as $a + (b + c)$ and this is succinctly written as $a + b + c$. Similarly, $(a \cdot b) \cdot c = a \cdot (b \cdot c) = a \cdot b \cdot c$.
This is *not true* for subtraction. $(a - b) - c$ is not the same as $a - (b - c)$. Indeed, the latter is $(a - b) + c$.
- **Multiplication distributes over addition and subtraction.** For any three numbers, $a \cdot (b + c)$ equals $a \cdot b + a \cdot c$. Similarly, $a \cdot (b - c) = a \cdot b - a \cdot c$.
- **Prime and Composite Numbers.** A positive number p is *prime* if the only numbers dividing it (that is, leaving 0 remainder) are 1 and p . Otherwise, the number is composite. The only exception to the rule is 1 which is neither prime nor composite. For instance, 17 is a prime, but $323 = 17 \times 19$ is not.
- **Floors and Ceilings.** Given any real number x , the *floor* $\lfloor x \rfloor$ is the *largest integer smaller than or equal to* x . So, $\lfloor 2 \rfloor = 2$, and $\lfloor 1.5 \rfloor = 1$, and $\lfloor -2.7 \rfloor = -3$. Similarly, the *ceiling* $\lceil x \rceil$ is the *smallest integer larger than or equal to* x . So, $\lceil 2 \rceil = 2$ and $\lceil 1.5 \rceil = 2$ and $\lceil -2.7 \rceil = -2$.
- **Exponentiating.** Given any positive integer n and any real x , the number x^n is a shorthand for $x \cdot x \cdot x \cdots x$, where x is multiplied with itself n times. So, $3^2 = 9$ and $(1.5)^2 = 2.25$ and $(-1)^3 = -1$.
The number x^0 is defined to be 1 for any x . For any negative number $y = -z$, we define $x^y = x^{-z} := 1/x^z$.

Some properties of exponentials: For any reals x, y and numbers a, b , we have

1. $x^{a+b} = x^a \cdot x^b$
2. $(xy)^a = x^a \cdot y^a$
3. $(x^a)^b = x^{ab}$

- **Logarithms.** The *logarithm* of a *positive* number a to the *positive base* $b \neq 1$, denoted as $\log_b a$ is the number ℓ such that $b^\ell = a$. Thus, $\log_3 81 = 4$ (since $3^4 = 81$) and $\log_{4/3}(16/9) = 2$ (since $(4/3)^2 = 16/9$), and $\log_{\sqrt{2}} 16 = 4$ (since $(\sqrt{2})^8 = 16$).

Some properties of logarithms: These properties follow by using the properties of exponential and the definition of logarithms

1. $\log_b 1 = 0$ for all $b > 0$.

2. $\log_b(xy) = \log_b x + \log_b y$ for all $b, x, y > 0$.
3. $\log_b(x^y) = y \log_b x$ for all $b, x, y > 0$.
4. $\log_b x = \frac{\log_c x}{\log_c b}$.

• **Basic Formulae.** You should be able to deduce the following

1. For any two numbers, $(a + b)^2 = a^2 + 2ab + b^2$
2. For any two numbers, $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$
3. For any three numbers, $(a + b + c)^2 = a^2 + b^2 + c^2 + 2(ab + bc + ca)$

• **Polynomials.** A polynomial is a formula of the form $p(x) = a_d x^d + a_{d-1} x^{d-1} + \dots + a_1 x + a_0$. x is called the *variable*. The *degree* of a polynomial is the largest power of x which participates in the polynomial. In the above example, this is d . The constants a_0, a_1, \dots, a_d are the *coefficients* of the polynomial.

For instance, $p(x) = x^2 + 2x + 1$ is a polynomial, and so is $q(x) = 5x^3 - 7x + 4$. The degree of $p(x)$ is 2 and its coefficients are 1, 2, 1, while the degree of $q(x)$ is 3 and its coefficients are 5, 0, -7, 4.

You should be able to solve these problems reasonably quickly. If not, come talk to me ASAP.

- What is the value of (a) $\lfloor 2.5 \rfloor + \lceil 3.75 \rceil$? (b) $(\lfloor \pi \rfloor)^{\lceil \pi \rceil}$?
- Are $1 + \lfloor x \rfloor$ and $\lfloor 1 + x \rfloor$ always equal? Are $\lceil \lfloor x \rfloor \rceil$ and $\lfloor \lceil x \rceil \rfloor$ always equal?
- If x and y are rational numbers, are $x + y$, $x - y$, and $x \cdot y$ always rational? How about x/y ?
- Which is bigger 3^{10} or 10^3 ?
- What is the value of (a) $\log_{1/8} 2$? (b) $\log_2 16$?
- Which is bigger, $\log_{10} 17$ or $\log_{17} 10$?