Math 38 - Graph Theory Maximum and perfect matchings

Nadia Lafrenière 04/27/2022

In a graph, a <u>matching</u> is a subgraph with maximal degree 1 (so every vertex is connected to at most one other vertex).

A maximal, but not maximum matching

M

A maximum matching, also a perfect matching




A vertex that appears in a matching is saturated, otherwise it is unsaturated.

A perfect matching in a graph is a matching that saturates every vertex.

Example

In the complete bipartite graph K_{mn} , there exists perfect matchings only if m=n. In this case, the matchings of graph K_{mn} represent bijections between two sets of size n. These are the permutations of n, so there are n! matchings.



- Perfect matchings can only occur when the number of vertices is even.
- That is not a sufficient condition, as shown by the claw.



No possible perfect matching, since the center vertex is saturated by any edge.

(2)Example Counting the perfect matchings in a complete graph. - K has no perfect matching if n is odd. - Otherwise, it has (n-1)(n-3)... 3.1 perfect matchings: - Label the vertices 1,...,n - Match vertex 1 with any of its neighbors; there are n-1 possible choices - As long as there are still unsaturated vertices, match the smallest unsaturated vertex with another one. The number of ways to do so is n-3, then n-5, ..., until there is only one way to do so. Maximum matchings A matching of a graph is maximal if no edge can be added. It is maximum if no other matching of this graph has more edges than it. Example Maximal Maximum Perfect => Maximum => Maximal In general, the converse is not true. Can we transform a maximal matching into a maximum matching? M-alternating path Let G be a graph and M be a matching of G. An M-alternating path is a path of G that alternates between edges in M and edges not in M.

a path of G that alternates between edges in M and edges not in M. An M-augmenting path is an M-alternating path with both endpoints unsaturated.




Remark: When M is maximum, there is no augmenting path.

Theorem (Berge, 1957) A matching M in a graph is a maximum matching if and only if the graph has no M-augmenting path. 3

<u>Proof</u> \Rightarrow follows from the remark above \Leftarrow We prove the converse: if it is not maximum, it has an augmenting path. If M is not maximum, then there is a matching M' with more edges.

We consider the subgraph H with edges that appear in exactly one of M and M' (not in both). Claim: components of H are all either even cycles or paths. - Even cycles have as many edges from M as from M'. This is because every endpoint can have at most one edge in M and one in M'. - Since IM'I>IMI, there is at least one path in H with more edges of M' than edges of M. This is a path that starts and ends with unsaturated vertices of M, so this is an M-augmenting path.

<u>Proof</u> of the claim: That means that every vertex of H has degree at most 2, and that cycles have even length. The maximal degree of H is 2 by construction. At most, one vertex can have one incident edge in M and one in M^{*}. If a cycle has odd length, then most edges belong to the same matching, and there must be two edges belonging to the same matching and incident to the same vertex. That contradicts the construction of a matching.



Matchings in bipartite graphs

Example: Job assignments If there are m jobs and n people, not all qualified for all the jobs, can we always fill all the jobs?



(The jobs cannot all be filled in this example).

Theorem (Hall's Theorem, 1935) Let G be a bipartite graph with the independent sets X and Y forming a partition of the vertices. G has a matching that saturates every vertex of X if and only if the neighborhood of every $S \subseteq X$ has order at least ISI.



Consequence: Stable marriages. Watch the video: https://www.numberphile.com/videos/stable-marriage-problem

Reference: Douglas B. West. Introduction to graph theory, 2nd edition, 2001. Section 3.1